

Datasheet for ABIN1302482

**anti-Bcl-2 antibody (AA 41-54)**

4 Images

4 Publications

[Go to Product page](#)

## Overview

Quantity:	0.1 mg
Target:	Bcl-2 (BCL2)
Binding Specificity:	AA 41-54
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Bcl-2 antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunoprecipitation (IP), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))

## Product Details

Immunogen:	Synthetic peptide corresponding to the amino acids 41-54 of human Bcl2
Clone:	Bcl-2-100
Isotype:	IgG1
Specificity:	The mouse monoclonal antibody Bcl-2/100 recognizes Bcl2, a 26 kDa intracellular protooncogen with anti-apoptotic effect, expressed in outer mitochondrial membrane, endoplasmic reticulum and nuclear envelope.
No Cross-Reactivity:	Mouse
Cross-Reactivity (Details):	Human



## Product Details

Purification: Purified by protein-A affinity chromatography.

Purity: > 95 % (by SDS-PAGE)

## Target Details

Target: Bcl-2 (BCL2)

Alternative Name: Bcl2 ([BCL2 Products](#))

Background: BCL2 apoptosis regulator, Bcl2 (B cell chronic lymphatic leukemia protein 2) is a proto-oncogen, which can contribute to tumorigenesis by counteracting apoptosis in various cell types. The anti-apoptotic effect of Bcl2 is performed by its interactions with suppressors and agonists of cell death and under physiological conditions it is regulated by proteolytic processing and phosphorylation. Bcl2 expression can be detected mainly in lymphoid tissues and in the basal cells of epithelial tissues. It is also a marker that can help in classification of lymphoproliferative diseases and in prognostics of some epithelial neoplasms., Bcl-2, PPP1R50

Gene ID: 596

UniProt: [P10415](#)

Pathways: [MAPK Signaling](#), [PI3K-Akt Signaling](#), [Apoptosis](#), [Caspase Cascade in Apoptosis](#), [Regulation of Muscle Cell Differentiation](#), [Cell-Cell Junction Organization](#), [Skeletal Muscle Fiber Development](#), [Autophagy](#), [Smooth Muscle Cell Migration](#), [Negative Regulation of intrinsic apoptotic Signaling](#)

## Application Details

Application Notes: Flow cytometry: Recommended dilution: 1-5 µg/mL. Intracellular staining.

Restrictions: For Research Use only

## Handling

Concentration: 1 mg/mL

Buffer: Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling Advice: **Do not freeze.**



## Handling


Storage: 4 °C

Storage Comment: Store at 2-8°C. Do not freeze.

## Publications

- Product cited in: Joubert, Marais, Maritz: "Influence of 2-methoxyestradiol on MCF-7 cells: an improved differential interference contrasting technique and Bcl-2 and Bax protein expression levels." in: **Biocell : official journal of the Sociedades Latinoamericanas de Microscopía Electronica ... et. al**, Vol. 33, Issue 1, pp. 67-70, (2009) ([PubMed](#)).
- Gugasyan, Christou, O'Reilly, Strasser, Gerondakis: "Bcl-2 transgene expression fails to prevent fatal hepatocyte apoptosis induced by endogenous TNFalpha in mice lacking RelA." in: **Cell death and differentiation**, Vol. 13, Issue 7, pp. 1235-7, (2006) ([PubMed](#)).
- Laflamme, Israël-Assayag, Cormier: "Apoptosis of bronchoalveolar lavage lymphocytes in hypersensitivity pneumonitis." in: **The European respiratory journal**, Vol. 21, Issue 2, pp. 225-31, (2003) ([PubMed](#)).
- Soilu-Hänninen, Ekert, Bucci, Syroid, Bartlett, Kilpatrick: "Nerve growth factor signaling through p75 induces apoptosis in Schwann cells via a Bcl-2-independent pathway." in: **The Journal of neuroscience : the official journal of the Society for Neuroscience**, Vol. 19, Issue 12, pp. 4828-38, (1999) ([PubMed](#)).


## Images



### Immunohistochemistry (Paraffin-embedded Sections)


**Image 1.** Immunohistochemistry staining of human placenta (paraffin-embedded sections) with anti-BCL2 (Bcl-2/100), 10 µg/mL.





#### Flow Cytometry

**Image 2.** Separation of human lymphocytes (red-filled) from neutrophil granulocytes (black-dashed) in flow cytometry analysis (intracellular staining) of human peripheral whole blood stained using anti-human BCL-2 (Bcl-2/100) purified antibody (concentration in sample 1  $\mu\text{g/mL}$ , GAM APC).



#### Flow Cytometry

**Image 3.** Flow cytometry intracellular staining pattern of human peripheral whole blood stained using anti-human BCL-2 (Bcl-2/100) purified antibody (concentration in sample 1  $\mu\text{g/mL}$ , GAM APC).

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN1302482.