antibodies -online.com





anti-IL3RA antibody

2 Images

2

Publications



Go to Product page

Overview

Quantity:	0.1 mg
Target:	IL3RA
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This IL3RA antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunoprecipitation (IP), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	IL3 receptor alpha chain expressed on the surface of transiently transfected COS cells
Clone:	6H6
Isotype:	lgG1
Specificity:	The mouse monoclonal antibody 6H6 recognizes an extracellular epitope of CD123 (interleukin 3 receptor alpha), a 60-70 kDa transmembrane protein expressed by myeloid precursors, megakaryocytes, macrophages, dendritic cells, mast cells, basophils, and some B cells. This antibody does not inhibit IL-3 binding to its receptor.
Cross-Reactivity (Details):	Human
Purification:	Purified by protein-A affinity chromatography.
Purity:	> 95 % (by SDS-PAGE)

Target Details

Target:	IL3RA
Alternative Name:	CD123 (IL3RA Products)
Background:	Interleukin 3 receptor subunit alpha,CD123 is the alpha chain of interleukin 3 receptor (IL-3R alpha). This subunit heterodimerizes with the interleukin 3 receptor beta chain (CD131), which is shared with other receptors. CD123 interacts with IL-3 specifically, but with low affinity, and association with the beta subunit confers high affinity binding to the receptor heterodimer. Both chains are required for signaling, but receptor activation and signal transduction depend on IL-3 binding to CD123 as the initial step.,IL3RA, MGC34174, hIL-3Ra
Gene ID:	3563
UniProt:	P26951
Pathways:	JAK-STAT Signaling

Application Details

Application Notes:	Flow cytometry: Recommended dilution: 1-4 μg/mL.
Restrictions:	For Research Use only
Handling	
Concentration:	1 mg/mL

Concentration:	1 mg/mL
Buffer:	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Do not freeze.
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Do not freeze.


Publications

Product cited in:

Martín-Gayo, Sierra-Filardi, Corbí, Toribio: "Plasmacytoid dendritic cells resident in human thymus drive natural Treg cell development." in: **Blood**, Vol. 115, Issue 26, pp. 5366-75, (2010) (PubMed).


Herling, Teitell, Shen, Medeiros, Jones: "TCL1 expression in plasmacytoid dendritic cells (DC2s) and the related CD4+ CD56+ blastic tumors of skin." in: **Blood**, Vol. 101, Issue 12, pp. 5007-9, (2003) (PubMed).

Images



Flow Cytometry

Image 1. Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD123 (6H6) purified antibody (concentration in sample 0,11 μ g/mL, GAM APC).



Flow Cytometry

Image 2. Separation of CD123 positive basophil granulocytes (red-filled) from neutrophil granulocytes (black-dashed) in flow cytometry analysis (surface staining) of peripheral whole blood stained using anti-human CD123 (6H6) purified antibody (concentration in sample 0,11 μ g/mL, GAM APC).