

Datasheet for ABIN135022
anti-BCL2L1 antibody (FITC)



[Go to Product page](#)

1 Image

1 Publication

Overview

Quantity:	0.1 mg
Target:	BCL2L1
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This BCL2L1 antibody is conjugated to FITC
Application:	Flow Cytometry (FACS)

Product Details

Immunogen:	Recombinant human Bcl-xS
Clone:	7B2-5
Isotype:	IgG3
Specificity:	Human/Mouse/Rat/Rhesus Bcl-xL
Characteristics:	Mouse Anti-Bcl-xL-FITC

Target Details

Target:	BCL2L1
Alternative Name:	Bcl-xL (BCL2L1 Products)
Background:	Apoptosis, or programmed cell death, is a well-documented phenomenon in many cellular systems. It plays a key role in tissue and organ development as well as in adult tissues during

Target Details

cell turnover. Apoptosis can be induced by a variety of internal and external stimuli including growth factor deprivation, cytokine treatment, antigen-receptor engagement, cell-cell interactions, irradiation and glucocorticoid treatment. Bcl-2 and one of its homologues, Bcl-xL, protect cells from apoptosis, while other homologues of Bcl-2 such as Bax, Bad and Bak have been shown to enhance apoptosis. Bcl-xL has been shown to block apoptosis which is induced by a variety of stimuli and, under certain conditions, offers greater protection against apoptosis than Bcl-2. In contrast, Bad and Bax inhibit the protective functions of Bcl-xL and Bcl-2, respectively. Although heterodimerization between Bcl-xL/Bad and Bcl-2/Bax was originally thought to be essential for the differential anti-apoptotic activity of Bcl-xL and Bcl-2, other results suggest that the formation of heterodimers may not be necessary for this death-repressing activity.

Pathways: [Apoptosis](#), [Negative Regulation of intrinsic apoptotic Signaling](#)

Application Details

Application Notes:

- **Applications:** FC - Quality tested , IHC-PS - Reported in literature , ICC - Reported in literature , WB - Reported in literature , IP - Reported in literature , ELISA - Reported in literature
- **Working Dilutions:** Flow Cytometry FITC and BIOT conjugates 3 g/106 cells PE conjugate 0.3 g/106 cells For flow cytometry, the suggested use of these reagents is in a final volume of 100 L Immunoblotting Purified (UNLB) antibody 1 g/mL

Sample Volume: 1 mL

Restrictions: For Research Use only

Handling

Concentration: 0.1 mg/mL

Buffer: 0.1 mg in 1.0 mL PBS/Sodium azide

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling Advice: **Protect conjugated products from light.**
Each reagent is stable for the period shown on the bottle label if stored as directed.

Storage: 4 °C

Storage Comment: Store at 2-8°C

Publications

Product cited in: Gottschalk, Boise, Oltvai, Accavitti, Korsmeyer, Quintáns, Thompson: "The ability of Bcl-x(L) and Bcl-2 to prevent apoptosis can be differentially regulated." in: **Cell death and differentiation**, Vol. 3, Issue 1, pp. 113-8, (2006) ([PubMed](#)).

Images




Image 1.