

Datasheet for ABIN135095

anti-CD3 epsilon antibody (Biotin)

2 Images



Overview

Quantity:	0.5 mg
Target:	CD3 epsilon (CD3E)
Reactivity:	Mouse
Host:	Rat
Clonality:	Monoclonal
Conjugate:	This CD3 epsilon antibody is conjugated to Biotin
Application:	Flow Cytometry (FACS)

Product Details

Immunogen:	IL-4 producing Th2 cell lines including D10
Clone:	C363-29B
Isotype:	lgG2b
Specificity:	Mouse CD3, Mr 25 kDa
Characteristics:	Rat Anti-Mouse CD3e-BIOT

Target Details

Target:	CD3 epsilon (CD3E)
Alternative Name:	CD3e (CD3E Products)
Background:	CD3, a member of the immunoglobulin superfamily of cell surface receptors, is comprised of
	five invariable chain ranging in size from 16-28 kDa and is closely associated with the T cell

	antigen receptor (TCR). It is expressed on all T cells of all mouse strains. CD3 plays a major role
	in signaling during antigen recognition, leading to T-cell activation.
Pathways:	TCR Signaling, CXCR4-mediated Signaling Events, Ubiquitin Proteasome Pathway
Application Details	
Application Notes:	 Applications: FC - Quality tested, IHC-FS - Reported in literature, IHC-PS - Reported in literature, IP - Reported in literature, Depletion - Reported in literature, Activ - Reported in literature, CMCD - Reported in literature Working Dilutions: Flow Cytometry FITC and BIOT conjugates 3 g/106 cells PE, APC, SPRD, and CY5 conjugates 1 g/106 cells For flow cytometry, the suggested use of these reagents is in a final volume of 100 L
Comment:	In vitro depletion of CD3+ cells, In vitro activation of T cells
Sample Volume:	1 mL
Restrictions:	For Research Use only
Handling	
Concentration:	0.5 mg/mL
Buffer:	0.5 mg in 1.0 mL of PBS/Sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Protect conjugated products from light.
	Each reagent is stable for the period shown on the bottle label if stored as directed.
Storage:	4°C
Storage Comment:	Store at 2-8°C




Image 1.

Flow Cytometry

Image 2. C57BL/6 mouse splenocytes were stained with Rat Anti-Mouse CD3ɛ-FITC.