

# Datasheet for ABIN135106 anti-CD5 antibody



Go to Product page

## 1 Image

Overview

Alternative Name:

Background:

Quantity:	0.5 mg
Target:	CD5
Reactivity:	Mouse
Host:	Rat
Clonality:	Monoclonal
Conjugate:	This CD5 antibody is un-conjugated
Application:	Flow Cytometry (FACS)
Product Details	
Immunogen:	Unknown
Immunogen: Clone:	Unknown B19-1
Clone:	B19-1
Clone: Isotype:	B19-1 IgG2a
Clone: Isotype: Specificity:	B19-1 IgG2a Mouse CD5, Mr 67 kDa
Clone:  Isotype:  Specificity:  Characteristics:	B19-1  IgG2a  Mouse CD5, Mr 67 kDa  Rat Anti-Mouse CD5-UNLB

CD5/Lyt-1 antigen is a monomeric type I transmembrane glycoprotein expressed on

CD5 (CD5 Products)

thymocytes, T lymphocytes, and a subset of B lymphocytes, but not on natural killer (NK) cells. It has been identified as the major ligand of the B-cell antigen CD72. The frequency of CD5+ B cells exhibits strain-dependent variation, and the phenotypic, anatomical, functional, developmental, and pathological characteristics of the CD5+ B cells suggest that they may represent a distinct lineage, known as B-1 cells. Binding of CD5 on the T cell surface can augment alloantigen- or mitogen-induced lymphocyte proliferation and induces increased cytosolic free calcium, IL-2 secretion, and IL-2R expression. It has been proposed that CD5 negatively regulates signal transduction mediated by the T-cell and B-cell receptors.

#### **Application Details**

#### Application Notes:

- Applications: FC Quality tested , IP Reported in literature , WB Reported in literature
- Working Dilutions: Flow Cytometry FITC and BIOT conjugates 1 g/106 cells PE, APC, and SPRD conjugates 0.2 g/106 cells For flow cytometry, the suggested use of these reagents is in a final volume of 100 L

Sample Volume:


1 mL

Restrictions:

For Research Use only

#### Handling

Concentration:	0.5 mg/mL
Buffer:	0.5 mg of purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. No preservatives or amine-containing buffer salts added
Preservative:	Without preservative
Handling Advice:	Each reagent is stable for the period shown on the bottle label if stored as directed.
Storage:	4 °C
Storage Comment:	Store at 2-8°C



### Flow Cytometry

**Image 1.** BALB/c mouse splenocytes were stained with Rat Anti-Mouse CD5-PE.