antibodies -online.com







anti-H-2Dk antibody





Overview

Quantity:	0.5 mg
Target:	H-2Dk
Reactivity:	Mouse
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This H-2Dk antibody is un-conjugated
Application:	Flow Cytometry (FACS)

Product Details

Immunogen:	C3H mouse splenocytes
Clone:	15-5-5S
Isotype:	lgG2a
Specificity:	Mouse H-2Dk
Characteristics:	Mouse Anti-Mouse H-2Dk-UNLB
Purification:	Purified

Target Details

Target:	H-2Dk
Abstract:	H-2Dk Products

Application Details


Application Notes:	 Applications: FC - Quality tested, IP - Reported in literature, CMCD - Reported in literature, ELISA - Reported in literature, Block - Reported in literature Working Dilutions: Flow Cytometry FITC, BIOT, and PE conjugates 1 g/106 cells For flow cytometry, the suggested use of these reagents is in a final volume of 100 L
Comment:	Complement-dependent cytotoxicity
Sample Volume:	1 mL
Restrictions:	For Research Use only
Handling	
Concentration:	0.5 mg/mL
Buffer:	0.5 mg of purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. No preservatives or amine-containing buffer salts added
Preservative:	Without preservative

Images

Storage:

Handling Advice:

Storage Comment:



4°C

Store at 2-8°C

Flow Cytometry

Each reagent is stable for the period shown on the bottle label if stored as directed.

Image 1. C3H/He mouse splenocytes were stained with Mouse Anti-Mouse H-2Dk-PE.