antibodies - online.com







anti-H-2Dk antibody (Biotin)





Overview

Quantity:	0.5 mg
Target:	H-2Dk
Reactivity:	Mouse
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This H-2Dk antibody is conjugated to Biotin
Application:	Flow Cytometry (FACS)

Product Details

Immunogen:	C3H mouse splenocytes
Clone:	15-5-5S
Isotype:	lgG2a
Specificity:	Mouse H-2Dk
Characteristics:	Mouse Anti-Mouse H-2Dk-BIOT

Target Details

Target:	H-2Dk	
Abstract:	H-2Dk Products	

Application Details

Application Details		
Application Notes:	 Applications: FC - Quality tested , IP - Reported in literature , CMCD - Reported in literature , ELISA - Reported in literature , Block - Reported in literature Working Dilutions: Flow Cytometry FITC, BIOT, and PE conjugates 1 g/106 cells For flow cytometry, the suggested use of these reagents is in a final volume of 100 L 	
Comment:	Complement-dependent cytotoxicity	
Sample Volume:	1 mL	
Restrictions:	For Research Use only	
Handling		
Concentration:	0.5 mg/mL	
Buffer:	0.5 mg in 1.0 mL of PBS/Sodium azide	
Preservative:	Sodium azide	

Handling Advice: Protect conjugated products from light.

should be handled by trained staff only.

Each reagent is stable for the period shown on the bottle label if stored as directed.


This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which

Storage: 4 °C

Storage Comment: Store at 2-8°C

Images

Precaution of Use:



Flow Cytometry

Image 1. C3H/He mouse splenocytes were stained with Mouse Anti-Mouse H-2Dk-PE.