antibodies -online.com





anti-Kyphoscoliosis Peptidase antibody (PE)

2 Images



Go to Product page

Overview

Quantity:	0.1 mg
Target:	Kyphoscoliosis Peptidase (KY)
Reactivity:	Pig
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Kyphoscoliosis Peptidase antibody is conjugated to PE
Application:	Flow Cytometry (FACS)

Product Details

Immunogen:	Fresh dd miniature swine thymocytes
Clone:	76-7-4
Isotype:	lgG2a
Specificity:	Porcine CD1, Mr 40 & 11 kDa
Characteristics:	Mouse Anti-Porcine CD1-PE

Target Details

Target:	Kyphoscoliosis Peptidase (KY)
Alternative Name:	CD1 (KY Products)
Background:	Porcine CD1 is a type I transmembrane glycoprotein and a member of the immunoglobulin
	superfamily of cell surface receptors. It has a domain organization similar to that of MHC class

Target Details

	I molecules and is expressed in association with 2-microglobulin. CD1 is found on B cells,
	macrophages and immature thymocytes. There is evidence for a role of CD1 in presentation of
	lipids and peptides to T cells.
Pathways:	Skeletal Muscle Fiber Development
Application Details	
Application Notes:	• Applications: FC - Quality tested , IHC-FS - Reported in literature , ICC - Reported in literature ,
	IP - Reported in literature , CMCD - Reported in literature
	• Working Dilutions: Flow Cytometry FITC and BIOT conjugates 1 g/106 cells PE conjugate 0.2
	g/106 cells For flow cytometry, the suggested use of these reagents is in a final volume of 100 L
Sample Volume:	1 mL
Restrictions:	For Research Use only
Handling	
Concentration:	0.1 mg/mL
Buffer:	0.1 mg in 1.0 mL of PBS/Sodium azide and a stabilizing agent
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Handling Advice:	Do not freeze!
	Protect conjugated products from light.
	Each reagent is stable for the period shown on the bottle label if stored as directed.
Storage:	4 °C
Storage Comment:	Store at 2-8°C




Image 1.

Flow Cytometry

Image 2. Porcine peripheral blood lymphocytes were stained with Mouse Anti-Porcine CD1-PE.