

Datasheet for ABIN135466

anti-CD3 epsilon antibody (Biotin)**2** Images[Go to Product page](#)

Overview

Quantity:	0.5 mg
Target:	CD3 epsilon (CD3E)
Reactivity:	Pig
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD3 epsilon antibody is conjugated to Biotin
Application:	Flow Cytometry (FACS)

Product Details

Immunogen:	Purified CD3 molecules from porcine thymus
Clone:	PPT3
Isotype:	IgG1
Specificity:	Porcine CD3, Mr 23 kDa
Characteristics:	Mouse Anti-Porcine CD3e-BIOT

Target Details

Target:	CD3 epsilon (CD3E)
Alternative Name:	CD3e (CD3E Products)
Background:	Porcine CD3 is a member of the T-cell receptor-associated CD3 complex. It is found on a subpopulation of thymocytes and on all pig T lymphocytes. MAb PPT3 is mitogenic when

Target Details

presented to peripheral blood mononuclear cells in immobilized form.

Pathways: [TCR Signaling, CXCR4-mediated Signaling Events, Ubiquitin Proteasome Pathway](#)

Application Details

Application Notes:	• Applications: FC - Quality tested , IHC-FS - Reported in literature , IHC-PS - Reported in literature , IP - Reported in literature , WB - Reported in literature , Sep - Reported in literature , Stim - Reported in literature , Cyto - Reported in literature • Working Dilutions: Flow Cytometry FITC, BIOT, and AF488 conjugates 1 g/106 cells PE, APC, and SPRD conjugates 0.2 g/106 cells For flow cytometry, the suggested use of these reagents is in a final volume of 100 L
Comment:	In vitro activation of T cells
Sample Volume:	1 mL
Restrictions:	For Research Use only

Handling

Concentration:	0.5 mg/mL
Buffer:	0.5 mg in 1.0 mL of PBS/Sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Protect conjugated products from light. Each reagent is stable for the period shown on the bottle label if stored as directed.
Storage:	4 °C
Storage Comment:	Store at 2-8°C




Image 1.

Flow Cytometry

Image 2. Porcine peripheral blood lymphocytes were stained with Mouse Anti-Porcine CD3ε-PE.