

Datasheet for ABIN135496 **anti-CD8 antibody**

2 Images



Overview

Quantity:	0.5 mg
Target:	CD8
Reactivity:	Cat
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD8 antibody is un-conjugated
Application:	Flow Cytometry (FACS)

Product Details

Immunogen:	Feline thymocytes
Clone:	FCD8
Isotype:	lgG1
Specificity:	Feline/Lion/Tsushima Leopard Cat CD8β, Mr 31 & 38 kDa
Characteristics:	Mouse Anti-Feline CD8-UNLB
Purification:	Purified

Target Details

Target:	CD8
Alternative Name:	CD8 (CD8 Products)
Background:	Feline CD8, a member of the immunoglobulin superfamily of cell surface receptors, is a type II

Target Details

transmembrane glycoprotein that is expressed as a heterodimer on the suppressor/cytotoxic subpopulation of peripheral T lymphocytes. It is present on approximately 63 % of thymocytes, 9 % of splenocytes, 20 % of lymph node cells, and 15 % of peripheral blood lymphocytes. CD8 functions as a co-receptor with MHC Class I-restricted T cell receptors in antigen recognition.

Application Details

Storage:

Storage Comment:

Application Details		
Application Notes:	 Applications: FC - Quality tested , IHC-FS - Reported in literature , ICC - Reported in literature , IP - Reported in literature , Sep - Reported in literature Working Dilutions: Flow Cytometry FITC and BIOT conjugates 1 g/106 cells PE conjugate 0.2 g/106 cells For flow cytometry, the suggested use of these reagents is in a final volume of 100 L 	
Sample Volume:	1 mL	
Restrictions:	For Research Use only	
Handling		
Concentration:	0.5 mg/mL	
Buffer:	0.5 mg of purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. No preservatives or amine-containing buffer salts added	
Preservative:	Without preservative	
Handling Advice:	Each reagent is stable for the period shown on the bottle label if stored as directed.	

4°C

Store at 2-8°C




Image 1.

ELISA

Image 2. Feline peripheral blood lymphocytes were stained with Mouse Anti-Feline CD8-PE.