



[Go to Product page](#)

Datasheet for ABIN1385060
anti-AAV VP1 antibody

1 Image

Overview

Quantity:	100 µL
Target:	AAV VP1
Reactivity:	Adeno-Associated Virus (AAV)
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This AAV VP1 antibody is un-conjugated
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), ELISA, Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from Adeno-Associated Virus 5 capsid protein VP1
Isotype:	IgG
Specificity:	This antibody will recognize many AAV strains, including AAV5, AAV4, AAV3B, AAV9, and AAV13
Cross-Reactivity:	Virus
Cross-Reactivity (Details):	AAV5
Purification:	Purified by Protein A.

Target Details

Target:	AAV VP1
---------	---------

Target Details


Alternative Name:	Adeno-Associated Virus Capsid Protein VP1 (AAV VP1 Products)
Target Type:	Viral Protein
Background:	<p>Synonyms: capsid protein Adeno-associated 5 virus, capsid protein [Adeno-associated virus - 5], capsid protein AAV5, Parvovirus coat protein VP1, capsid protein.</p> <p>Background: Capsid protein self-assembles to form an icosahedral capsid with a T=1 symmetry, about 22 nm in diameter, and consisting of 60 copies of three size variants of the capsid protein VP1, VP2 and VP3 which differ in their N-terminus. The capsid encapsulates the genomic ssDNA. Binds to host cell heparan sulfate and uses host ITGA5-ITGB1 as coreceptor on the cell surface to provide virion attachment to target cell. This attachment induces virion internalization predominantly through clathrin-dependent endocytosis. Binding to the host receptor also induces capsid rearrangements leading to surface exposure of VP1 N-terminus, specifically its phospholipase A2-like region and putative nuclear localization signal(s). VP1 N-terminus might serve as a lipolytic enzyme to breach the endosomal membrane during entry into host cell and might contribute to virus transport to the nucleus</p>

Application Details

Application Notes:	WB 1:100-1000 IHC-P 1:100-500 IF(IHC-P) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months



ELISA

Image 1. Antigen: 0.2ug/100ul Primary: Antiserum, 1:500, 1:1000, 1:2000, 1:4000, 1:8000, 1:16000, 1:32000, Secondary: HRP conjugated Goat Anti-Rabbit IgG -HRP) at 1:5000, TMB staining, Read the data in MicroplateReader by 450nm.