
# antibodies -online.com





## anti-FAM92A1 antibody (AA 101-200)



## Overview

Quantity:	100 μL
Target:	FAM92A1
Binding Specificity:	AA 101-200
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FAM92A1 antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human FAM92A1
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Rabbit
Purification:	Purified by Protein A.

## **Target Details**

Target:	FAM92A1
Alternative Name:	FAM92A1 (FAM92A1 Products)

#### Target Details

Bac	kar	ound:

Synonyms: Family with sequence similarity 92, member A1, FLJ38979, Hypothetical protein LOC137392, Protein FAM92A1, F92A1\_HUMAN.

Background: Made up of nearly 146 million bases, chromosome 8 encodes about 800 genes. Translocation of portions of chromosome 8 with amplifications of the c-Myc gene are found in some leukemias and lymphomas, and typically associated with a poor prognosis. Portions of chromosome 8 have been linked to schizophrenia and bipolar disorder. Trisomy 8, also known as Warkany syndrome 2, most often results in early miscarriage but is occasionally seen in a mosaic form in surviving patients who suffer to a varying degree from a number of symptoms including retarded mental and motor development, and certain facial and developmental defects. WRN is a DNA helicase encoded by chromosome 8 and shown defective in those with the early aging disorder Werner syndrome. Chromosome 8 is also associated with Pfeiffer syndrome, congenital hypothyroidism and Waardenburg syndrome. The FAM92A1 gene product has been provisionally designated FAM92A1 pending further characterization.

Gene ID:

137392

#### **Application Details**

Application Notes:

WB 1:300-5000

ELISA 1:500-1000

IHC-P 1:200-400

IHC-F 1:100-500

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

ICC 1:100-500

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

## Handling

Storage:	4 °C,-20 °C	
Storage Comment:	t: Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	
Expiry Date:	12 months	