

Datasheet for ABIN1385154  
**anti-MDGA2 antibody (AA 751-850)**



[Go to Product page](#)

2 Images

## Overview

Quantity:	100 µL
Target:	MDGA2
Binding Specificity:	AA 751-850
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MDGA2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC)

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human MDGA2
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse,Dog,Cow,Sheep,Horse,Chicken
Purification:	Purified by Protein A.

## Target Details

Target:	MDGA2
---------	-------

## Target Details

---

Alternative Name:	MDGA2 ( <a href="#">MDGA2 Products</a> )
Background:	Synonyms: MAMDC1, c14_5286, MAM domain-containing glycosylphosphatidylinositol anchor protein 2, MAM domain-containing protein 1, MDGA2, UNQ8188/PRO23197 Background: May be involved in cell-cell interactions.
Gene ID:	161357
UniProt:	<a href="#">Q7Z553</a>

## Application Details


---

Application Notes:	WB 1:300-5000 ELISA 1:500-1000 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 ICC 1:100-500
Restrictions:	For Research Use only

## Handling


---

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months



#### Immunohistochemistry (Paraffin-embedded Sections)

**Image 1.** Formalin-fixed and paraffin embedded human glioma labeled with Rabbit Anti-MDGA2 Polyclonal Antibody, Unconjugated at 1:200 followed by conjugation to the secondary antibody and DAB staining



#### Immunohistochemistry (Paraffin-embedded Sections)

**Image 2.** Formalin-fixed and paraffin embedded rat brain labeled with Rabbit Anti-MDGA2 Polyclonal Antibody, Unconjugated at 1:200 followed by conjugation to the secondary antibody and DAB staining