

Datasheet for ABIN1385163 anti-Lamin B2 antibody (AA 61-160)



Overview

Quantity:	100 μL
Target:	Lamin B2 (LMNB2)
Binding Specificity:	AA 61-160
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Lamin B2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Lamin B2
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Cow,Sheep
Purification:	Purified by Protein A.

Target Details

Target: Lamin B2 (LMNB2)

Target Details

Alternative Name:	Lamin B2 (LMNB2 Products)
Background:	Synonyms: Alternative namesLAMB 2, LAMB2, Lamin-B2, LMN 2, LMN B2, LMN2, LMNB 2,
	LMNB2, LMNB2_HUMAN, MGC2721.
	Background: A unique family of Cysteine proteases has been described that differs in sequence
	structure and substrate specificity from any previously described protease family. This family,
	termed CED-3/ICE, functions as key components of the apoptotic machinery and act to destroy
	specific target proteins which are critical to cellular longevity. Nuclear lamins are critical to
	maintaining the integrity of the nuclear envelope and cellular morphology as components of the
	nuclear lamina, a fibrous layer on the nucleoplasmic side of the inner nuclear membrane, which
	is thought to provide a framework for the nuclear envelope and may also interact with
	chromatin. B-type lamins undergo a series of modifications, such as farnesylation and
	phosphorylation. Increased phosphorylation of the lamins occurs before envelope
	disintegration and probably plays a role in regulating lamin associations. Nuclear Lamin B is
	fragmented as a consequence of apoptosis by an unidentified member of the ICE family.
Gene ID:	84823
UniProt:	Q03252
Pathways:	Apoptosis, Caspase Cascade in Apoptosis
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL

Handling

Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months