# antibodies -online.com





# anti-PCDHGA9 antibody (AA 351-450)



Go to Product page

$\sim$	
( )\/△	rview
$\cup$	1 410 44

Quantity:	100 μL
Target:	PCDHGA9
Binding Specificity:	AA 351-450
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PCDHGA9 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc)), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC)

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human PCDHGA9
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat
Purification:	Purified by Protein A.

## **Target Details**

Target:	PCDHGA9
Alternative Name:	PCDHGA9 (PCDHGA9 Products)

#### **Target Details**

Buffer:

Preservative:

Precaution of Use:

ProClin

handled by trained staff only.

Synonyms: PCDH-gamma-A9, Protocadherin gamma A9, PCDG9_HUMAN.  Background: Protocadherins are a large family of cadherin-like cell adhesion proteins that are
Rackground: Protocodharing are a large family of codharin like call adhasian protoins that are
background. Protocaunerins are a large family of caunemininke cell aunesion proteins that are
involved in the establishment and maintenance of neuronal connections in the brain. There are
three protocadherin (PCDH) gene clusters, designated alpha, beta and gamma, all of which
contain multiple tandemly arranged genes. PCDHGA9 (Protocadherin gamma-A9) is a 932
amino acid that is one of 22 proteins encoded by the protocadherin gamma cluster. The
protocadherein gamma cluster consists of three subfamilies (A, B and C) and PCDHGA9 is a
member of the gamma subfamily A. PCDHGA9 is a type I transmembrane receptor containing
six cadherin motifs and is expressed in the central nervous system where it localizes to
synapses. Members of the gamma cluster of protocadherins are essential for neuronal survival.
There are two isoforms of PCDHGA9 that are produced as a result of alternative splicing
events.
56107
WB 1:300-5000
ELISA 1:500-1000
IHC-P 1:200-400
IHC-F 1:100-500
IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200
ICC 1:100-500
For Research Use only
Liquid
1 μg/μL

0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

## Handling

Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months