
# antibodies -online.com







#### Overview

Quantity:	100 μL
Target:	ube3a
Binding Specificity:	AA 701-800
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ube3a antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human UBE3A/E6-AP
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.
Target Details	

#### Target Details

Target: ube3a

#### **Target Details**

UBE3A/E6-AP (ube3a Products) Alternative Name: Background: Synonyms: ANCR, Angelman syndrome, AS, CTCL tumor antigen se37 2, E6 AP, E6AP antibody E6AP ubiquitin protein ligase, E6AP ubiquitin protein ligase, E6AP ubiquitin-protein ligase, EPVE6AP, HPVE6A, Human papilloma virus E6 associated protein Angelman syndrome, Human papilloma virus E6 associated protein, Human papillomavirus E6-associated protein, NY REN 54 antigen, NY REN 54 antigen, Oncogenic protein associated protein E6 AP, Oncogenic protein associated protein E6AP, Oncogenic protein-associated protein E6-AP, Renal carcinoma antigen NY REN 54, Renal carcinoma antigen NY-REN-54, UBE 3A, Ube3a, UBE3A protein, UBE3A\_HUMAN, Ubiquitin protein ligase E3A, Ubiquitin-protein ligase E3A. Background: E6-associating protein is a component of the ubiquitin-mediated proteolytic pathway, which selectively targets proteins for degradation by the 26S proteasome. Ubiquitin (Ub) is directly conjugated to protein substrates by the transfer of Ub from an E2 ubiquitin conjugating enzyme to the target protein. This conjugation is facilitated by the enzymatic activity of E3 ubiquitin ligase family members such as E6-AP. Several substrates of E6-AP have been identified and include the tumor suppressor protein p53 and the mammalian homolog of Rad23, HHR23A. Previous studies have indicated that E6-AP associates with the human papillomavirus E6 oncogene, which complexes with p53 and thereby potentiates E6-AP mediated ubiquitination of p53. Genetic mutations that impair E6-AP activity result in the accumulation of p53 in the cytoplasm, and, in many instances, these mutations are associated with the development of the rare neurodevelopmental disorder Angelman syndrome (AS), which is characterized by severe motor dysfunction and mental retardation. Gene ID: 7337 Pathways: Intracellular Steroid Hormone Receptor Signaling Pathway **Application Details** Application Notes: ELISA 1:500-1000 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 ICC 1:100-500 Restrictions: For Research Use only

## Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months