antibodies -online.com





anti-ZNF239 antibody (AA 275-350)



Go to Product page

()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL	
Target:	ZNF239	
Binding Specificity:	AA 275-350	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ZNF239 antibody is un-conjugated	
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ZNF239
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Cow,Sheep,Pig,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	ZNF239
Alternative Name:	ZNF239 (ZNF239 Products)

Target Details

Background:

Synonyms: HOK 2, Kruppel zinc finger protein 2, MOK 2, MOK2, OTTHUMP00000019484, OTTMUSP00000024957, zfp 239, Zinc finger protein C2H2 homologous to mouse MOK 2, zinc finger protein 239, Zinc finger protein HOK 2, Zinc finger protein HOK-2, zinc finger protein MOK 2, Zinc finger protein MOK-2, ZN239_HUMAN, Znf239.

Background: Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The

Background: Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Kr_ppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZNF239, also known as Zinc finger protein MOK-2 or HOK-2, is a 458 amino acid protein belonging to the Kr_ppel C2H2-type zinc-finger protein family. Localized to the nucleus, ZNF239 contains nine C2H2-type zinc finger domains. Due to the presence of these domains, ZNF239 may be involved in transcriptional regulation. ZNF239 is present at high levels in brain, breast and testis, and has no expression in liver or placenta.

Application Details

Application Notes:	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C

Handling

Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months