antibodies -online.com





Datasheet for ABIN1385329

anti-MEPE antibody (AA 201-300)



Overview

Quantity:	100 μL
Target:	MEPE
Binding Specificity:	AA 201-300
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MEPE antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunocytochemistry (ICC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human MEPE
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human
Purification:	Purified by Protein A.

Target Details

|--|

Target Details

Alternative Name:	MEPE (MEPE Products)
Background:	Synonyms: Matrix extracellular phosphoglycoprotein, MEPE, MEPE_HUMAN, 0F45,
	Osteoblast/osteocyte factor 45.
	Background: MEPE is a 525 amino acid extracellular matrix protein. Expressed in osteocytes
	and brain, MEPE is a regulator of bone metabolism that is thought to mediate mineralization
	and demineralization within the osteocyte microenvironment. MEPE contains an RGD cell-
	attachment motif and shares molecular similarities with several dentin-bone extracellular
	matrix RGD-containing phosphoglycoproteins, including OPN (osteopontin) and DSP (dentin
	sialophosphoprotein). Via its ability to control bone mineralization, MEPE is associated with
	various developmental events such as skeletogenesis, bone regeneration and odontogenesis.
	MEPE is secreted in hypophosphatemic osteomalacia tumors, suggesting a possible role in the
	pathophysiology of bone-related cancers.
Gene ID:	56955
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months