.-online.com antibodies

Datasheet for ABIN1385379 anti-RNF150 antibody

Image



Overview

Quantity:	100 µL
Target:	RNF150
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RNF150 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human RNF150
Isotype:	lgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.


Target Details

Target:	RNF150
Alternative Name:	RNF150 (RNF150 Products)
Background:	Synonyms: KIAA1214, MGC125502, Ring finger protein 150, RNF 150, RN150_HUMAN. Background: RNF150 contains a RING-type zinc finger, a motif known to be involved in protein-
	protein and protein-DNA interactions. The specific function of this protein is unknown.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1385379 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

Target Details	
Gene ID:	57484
Application Details	
Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Paraformaldehyde-fixed, paraffin embedded rat brain tissue, Antigen retrieval by boiling in sodium citrate buffer(pH6) for 15min, Block endogenous peroxidase by 3% hydrogen peroxide for 30 minutes, Blocking buffer (normal goat serum) at 37°C for 20min, Antibody incubation with Rabbit Anti-RNF150 Polyclonal Antibody, Unconjugated at 1:400 overnight at 4°C, followed by a conjugated secondary and DAB staining