



[Go to Product page](#)

Datasheet for ABIN1385388
anti-MAP9 antibody

Overview

Quantity:	100 µL
Target:	MAP9
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MAP9 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human MAP-9
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	MAP9
Alternative Name:	MAP-9 (MAP9 Products)
Background:	Synonyms: ASAP, Aster associated protein, MAP9, MAP 9, microtubule associated protein 9, MAP9_HUMAN. Background: Microtubules, the primary component of the cytoskeletal network, interact with

Target Details

proteins called microtubule-associated proteins (MAPs).MAP9 is a microtubule-associated protein required for spindle function, mitotic progression, and cytokinesis.The microtubule-associated proteins can be divided into two groups, structural and dynamic. The MAP proteins function to stimulate tubulin assembly, enhance microtubule stability, influence the spatial distribution of microtubules within cells and utilize microtubule polarity to translocate cellular components. MAP-9 (microtubule-associated protein 9), also known as ASAP, is a 647 amino acid cytoplasmic protein that is constitutively expressed during the cell cycle. MAP-9 localizes to microtubules in interphase, associates with the mitotic spindle during mitosis and localizes to the central body during cytokinesis. Involved in organization of the bipolar mitotic spindle, MAP-9 is required for bipolar spindle assembly, mitosis progression and cytokinesis. MAP-9 may be involved in stabilizing interphase microtubules. Two isoforms of MAP-9 are produced due to alternative splicing events.

Gene ID: 79884

Application Details

Application Notes: WB 1:300-5000
IHC-P 1:200-400
IF(IHC-P) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: 4 °C,-20 °C

Storage Comment: Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Expiry Date: 12 months