antibodies - online.com







anti-Nucleolar Protein 1 (NOL1) antibody



Image



Overview

Quantity:	100 μL
Target:	Nucleolar Protein 1 (NOL1)
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	Un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details


Immunogen:	KLH conjugated synthetic peptide derived from human NOL1
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	Nucleolar Protein 1 (NOL1)
Alternative Name:	NOL1 (NOL1 Products)
Background:	Synonyms: proliferation Associated Nucleolar Antigen: NOL 1, NUCLEOLAR ANTIGEN p120, NOP2_HUMAN.
	Background: Proliferation-associated nucleolar antigen is expressed in mid G1 phase with peak

Target Details

	level during the S phase and a rapid degradation during late mitosis. Its expression in breast
	carcinoma is correlated with patient prognosis.
Gene ID:	4839
Application Details	
Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Paraformaldehyde-fixed, paraffin embedded Mouse placenta, Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min, Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes, Blocking buffer (normal goat serum) at 37°C for 30min, Antibody incubation with NOP2 Polyclonal Antibody, Unconjugated at 1:400 overnight at 4°C, followed by a conjugated secondary antibody for 20 minutes and DAB staining.