antibodies -online.com





anti-FIG4 antibody (AA 1-100)



Overview

Quantity:	100 μL
Target:	FIG4
Binding Specificity:	AA 1-100
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FIG4 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FIG4
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human, Mouse, Dog, Sheep, Pig, Chicken, Monkey
Purification:	Purified by Protein A.

Target Details

Target: FIG4

Target Details

Alternative Name:	FIG4/SAC3 (FIG4 Products)
Background:	Synonyms: 5-bisphosphate 5-phosphatase, Fig4, FIG4_HUMAN, KIAA0274, ALS11,
	Phosphatidylinositol 3, hSac3, Phosphatidylinositol 3,5 bisphosphate 5 phosphatase,
	Polyphosphoinositide phosphatase, SAC domain containing protein 3, SAC domain-containing protein 3, SAC3.
	Background: FIG4 is a 907 amino acid protein that contains one SAC domain through which it is
	thought to function as a phosphoinositide phosphatase that may play an important role in
	signal transduction and vesicle trafficking. Defects in the gene encoding FIG4 are the cause of
	Charcot-Marie-Tooth disease type 4J (CMT4J) and amyotrophic lateral sclerosis type 11
	(ALS11), both of which negatively affect the nervous system. CMT4J is a recessive
	demyelinating disorder of the peripheral nervous system and is characterized by reduced motor
	nerve conduction velocities and axonal degeneration. Unlike CMT4J, ALS11 is characterized by
	the degeneration of upper motor neurons in the brain and lower neurons in the spinal cord,
	causing paralysis and, ultimately, death.
Pathways:	Inositol Metabolic Process
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months