

[Go to Product page](#)

Datasheet for ABIN1385666 **anti-MEF2B antibody**

Overview

Quantity:	100 µL
Target:	MEF2B
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MEF2B antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human MEF2B
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	MEF2B
Alternative Name:	MEF2B (MEF2B Products)
Background:	Synonyms: PRDM3, MADS box transcription enhancer factor 2, MDS1 EVI1, Mef2b, MEF2B_HUMAN, MEF-2B, myocyte enhancer factor 2B, myocyte specic enhancer factor 2B, Myocyte-specific enhancer factor 2B, RSRFR2, serum response factor like protein 2, Serum

Target Details

response factor-like protein 2, XMEF2.

Background: The myocyte enhancer factor-2 (MEF-2) family of transcription factors associate with co-repressors or co-activators to regulate development and function of T cells, neuronal cells, and muscle cells. Four family members, termed MEF-2A, -2B, -2C, and -2D, arise from alternatively spliced transcripts. These members bind as homo- and heterodimers to the MEF-2 site in the promoter region of affected genes. Differential regulation in the expression of the four transcripts implies functional distinction for each during embryogenesis and development. The process of differentiation from mesodermal precursor cells to myoblasts has led to the discovery of a variety of tissue-specific factors that regulate muscle gene expression. The myogenic basic helix-loop-helix proteins, including MyoD, myogenin, Myf-5, and MRF4, are one class of identified factors. The MEF-2 family represents a second class of DNA binding regulatory proteins. Each of these proteins binds to the MEF-2 target DNA sequence present in the regulatory regions of many muscle-specific genes.

Application Details

Application Notes: WB 1:300-5000
IHC-P 1:200-400
IF(IHC-P) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: 4 °C,-20 °C

Storage Comment: Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Expiry Date: 12 months