.-online.com antibodies

Datasheet for ABIN1385668 anti-PDE4DIP antibody



Overview

Quantity:	100 µL
Target:	PDE4DIP
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PDE4DIP antibody is un-conjugated
Application:	Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin- embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CMYA2/PDE4DIP
Isotype:	lgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	PDE4DIP
Alternative Name:	CMYA2 (PDE4DIP Products)
Background:	Synonyms: Cardiomyopathy associated protein 2, Cardiomyopathy-associated protein 2, CMYA2, MMGL, MYOME_HUMAN, Myomegalin, Pde4dip, Phosphodiesterase 4D interacting
	protein, Phosphodiesterase 4D-interacting protein.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1385668 | 03/06/2024 | Copyright antibodies-online. All rights reserved.

	Background: May function as an anchor sequestering components of the cAMP-dependent
	pathway to Golgi and/or centrosomes.Myomegalin, is a 2,346 amino acid protein that contains
	one NBPF domain and localizes to the nucleus, cytoplasm, centrosome and Golgi apparatus.
	Expressed at high levels in fetal and adult heart and at lower levels in brain and placenta,
	myomegalin is thought to function as an anchoring protein that sequesters members of the
	cAMP-dependent pathway to the Golgi and to centrosomes, thereby mediating cAMP pathway
	dynamics. Translocations in the gene that encodes myomegalin are associated with
	myeloproliferative disorders (MBDs), a group of diseases caused by an overproduction of blood
	cells. Myomegalin exists as twelve isoforms due to alternative splicing events.
Gene ID:	9659
Pathways:	SARS-CoV-2 Protein Interactome
Application Details	
Application Notes:	IHC-P 1:200-400
	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months