# antibodies -online.com







$\sim$			
	N/P	r\/	i⊢₩

Quantity:	100 μL	
Target:	CTDSPL	
Binding Specificity:	AA 175-276	
Reactivity:	Human, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CTDSPL antibody is un-conjugated	
Application:	Western Blotting (WB), Flow Cytometry (FACS), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human CTDSPL
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Cow,Sheep,Pig,Horse,Chicken
Purification:	Purified by Protein A.

#### **Target Details**

Target: CTDSPL

### **Target Details**

g		
Alternative Name:	Ctdspl (CTDSPL Products)	
Background:	Synonyms: Carboxy-terminal domain RNA polymerase II polypeptide A small phosphatase 3,	
	CTD small phosphatase-like protein, CTDSL_HUMAN, CTDSP-like, Ctdspl, hYA22, N-like protein,	
	NLI-interacting factor 1, Nuclear LIM interactor-interacting factor 1, Protein YA22, RBSP3, SCP3	
	Small C-terminal domain phosphatase 3, Small CTD phosphatase 3.	
	Background: Preferentially catalyzes the dephosphorylation of 'Ser-5' within the tandem 7	
	residues repeats in the C-terminal domain (CTD) of the largest RNA polymerase II subunit	
	POLR2A. Negatively regulates RNA polymerase II transcription, possibly by controlling the	
	transition from initiation/capping to processive transcript elongation (By similarity). Recruited	
	by REST to neuronal genes that contain RE-1 elements, leading to neuronal gene silencing in	
	non-neuronal cells.	
Gene ID:	10217	
Application Details		
Application Notes:	WB 1:300-5000	
	ELISA 1:500-1000	
	FCM 1:20-100	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	

## Handling

Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months