antibodies -online.com







anti-PBLD1 antibody (AA 60-110)

Images



Publication



| <i>ا</i> ۱ | 111 | SKI | / 1 | ~ 1 A |
|------------|-------|------|-------------|-------|
| u | \ / F | ٠١ ' | \/ | -1 |
| \sim | V (| ~ I | VΙ | ev |

| Quantity: | 100 μL | |
|----------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------|--|
| Target: | PBLD1 | |
| Binding Specificity: | AA 60-110 | |
| Reactivity: | Human, Mouse, Rat | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This PBLD1 antibody is un-conjugated | |
| Application: | Western Blotting (WB), ELISA, Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human PBLD/MAWBP | |
|-------------------|----------------------------------------------------------------|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Human, Mouse, Rat | |
| Purification: | Purified by Protein A. | |

Target Details

| Target: | PBLD1 |
|---------------------------------------------------------------------------------------------|-----------------------|
| Alternative Name: | PBLD (PBLD1 Products) |
| Background: Synonyms: MAWBP, MAWDBP, Phenazine biosynthesis-like domain-containing protein, | |

binding protein, Unknown protein 32 from 2D-page of liver tissue, PBLD Background: MAWDBP (MAWD binding protein), also known as PBLD (phenazine biosynthesis-like protein domain containing) or MAWBP, is a 288 amino acid protein that belongs to the phenazine biosynthesis-like protein (PhzF) family. It has been suggested that MAWDBP is the only representative of the PhzF family in the human genome. Expressed in most tissues, MAWDBP is a WD-40 repeat-containing _-propeller protein believed to participate in the MAPK signaling pathway. Involved in multiple basic cellular functions, expression of MAWDBP is elevated in several disease processes, including insulin resistance, folate deficiency and hypotension. It is thought that MAWDBP may also be involved in carcinogenesis.

Gene ID: 64081

UniProt: P30039

Application Details

Application Notes: WB 1:300-5000

ELISA 1:500-1000 IHC-P 1:200-400

IF(IHC-P) 1:50-200

Restrictions: For Research Use only

Handling


| Format: | Liquid | |
|--------------------|--------------------------------------------------------------------------------------------------------------------|--|
| Concentration: | 1 μg/μL | |
| Buffer: | 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. | |
| Storage: | 4 °C,-20 °C | |
| Storage Comment: | Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. | |
| Expiry Date: | 12 months | |

Product cited in:

Liu, Zhao, Lu, Fan, Wang: "Proteomic study on usnic-acid-induced hepatotoxicity in rats." in:


Journal of agricultural and food chemistry, Vol. 60, Issue 29, pp. 7312-7, (2012) (PubMed).

Images



ELISA

Image 1. Antigen: 0.2ug/100ul, Primary: Antiserum, 1:500, 1:1000, 1:2000, 1:4000, 1:8000, 1:16000, 1:32000, Secondary: HRP conjugated Goat-Anti-Rabbit IgG at 1: 5000, TMB staining, Read the data in MicroplateReader by 450nm



ELISA

Image 2. Antigen: 2 μ g/100 μ L Primary: Antiserum, 1:500, 1:1000, 1:2000, 1:4000, 1:8000, 1:16000, 1:32000; Secondary: HRP conjugated Rabbit Anti-Goat IgG at 1: 5000; TMB staining Read the data in Microplate Reader by 450nm.