# antibodies -online.com





## anti-NOS1AP antibody (AA 128-170)



## Overview

Quantity:	100 μL
Target:	NOS1AP
Binding Specificity:	AA 128-170
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NOS1AP antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)),
	Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunocytochemistry (ICC),
	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry
	(Frozen Sections) (IHC (fro))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human CAPON
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Cow,Sheep,Horse
Purification:	Purified by Protein A.

## **Target Details**

Target:	NOS1AP
Alternative Name:	CAPON (NOS1AP Products)

#### **Target Details**

#### Background:

Synonyms: NOS1AP, C terminal PDZ domain ligand of neuronal nitric oxide synthase CAPON, C terminal PDZ domain ligand of neuronal nitric oxide synthase, C terminal PDZ ligand of neuronal nitric oxide synthase protein, C-terminal PDZ ligand of neuronal nitric oxide synthase protein, CAPON, CAPON\_HUMAN, Carboxyl terminal PDZ ligand of neuronal nitric oxide synthase protein, Carboxyl-terminal PDZ ligand of neuronal nitric oxide synthase protein, Ligand of neuronal nitric oxide synthase with carboxyl terminal PDZ domain, MGC138500, Nitric oxide synthase 1 neuronal adaptor protein, Nitric oxide synthase 1 adaptor protein. Background: CAPON (carboxy-terminal PDZ ligand of nNOS) selectively binds within the 100 amino acid PDZ domain of the neuronal nitric oxide synthase (nNOS), but not to endothelial NOS or inducible NOS, and sequesters nNOS in the cytosol. Biosynthesis of the neurotransmitter nitric oxide (NO) requires the association of nNOS with various synaptic proteins, including syntrophin, postsynaptic density (PSD)95 and PSD93 through a scaffolding PDZ domain. These proteins facilitate the transport of nNOS to the plasma membrane, where it is catalytically activated by NMDA-receptor mediated calcium channels. The association of nNOS with PSD95 or PSD93 is regulated by CAPON. The carboxy terminus of CAPON binds to the PDZ domain, competes with PSD95 and PSD93 for binding to nNOS and in turn prevents the translocation and catalytic activation of nNOS.

### **Application Details**

Buffer:

Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL

0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.

## Handling

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months