

Datasheet for ABIN1385801
anti-NSF antibody (AA 151-250)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	NSF
Binding Specificity:	AA 151-250
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NSF antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunocytochemistry (ICC), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human NSF
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit,Zebrafish
Purification:	Purified by Protein A.

Target Details

Target:	NSF
---------	-----

Target Details

Alternative Name:	NSF (NSF Products)
Background:	<p>Synonyms: N ethylmaleimide sensitive factor, N ethylmaleimide sensitive factor, N ethylmaleimide sensitive factor like protein, N ethylmaleimide sensitive fusion protein, N ethylmaleimide sensitive fusion protein, N-ethylmaleimide-sensitive fusion protein, NEM sensitive fusion protein, NEM sensitive fusion protein, NEM-sensitive fusion protein, NSF, NSF_HUMAN, SKD 2, SKD2, SKD2, Vesicle fusing ATPase, Vesicle fusing ATPase, Vesicle-fusing ATPase, Vesicular fusion protein NSF, Vesicular fusion protein NSF, Vesicular-fusion protein NSF.</p> <p>Background: Several protein-protein interactions are essential to membrane fusion during endocytosis. Membrane fusion requires interaction among SNARE1 proteins associated with both donor and acceptor membranes (1,2). Following membrane fusion, the -SNAP cytoplasmic adapter protein binds to the SNARE complex. N-ethylmaleimide-sensitive factor (NSF), a hexameric ATPase, then associates with the -SNAP/SNARE complex to mediate SNARE disassembly during membrane fusion (3,4). The ATPase activity of NSF induces a conformational change in the -SNAP/SNARE complex that leads to its dissociation from the membrane, membrane fusion and eventual recycling of the SNARE complex for subsequent membrane fusion (3,4).</p>

Application Details

Application Notes:	WB 1:300-5000 ELISA 1:500-1000 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 ICC 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.

Handling

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months