antibodies - online.com







anti-SFRS7 antibody (AA 2-70)





_						
0	V	0	r٧	/[Θ	M

Quantity:	100 μL
Target:	SFRS7
Binding Specificity:	AA 2-70
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SFRS7 antibody is un-conjugated
Application:	ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffinembedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details


Immunogen:	KLH conjugated synthetic peptide derived from human SFRS7	
Isotype:	IgG	
Cross-Reactivity:	Rat	
Predicted Reactivity:	Human,Mouse,Cow,Chicken,Zebrafish	
Purification:	Purified by Protein A.	
Target Details		

Target Details

|--|

Target Details

SFRS7/9G8/SRp20 (SFRS7 Products)		
Synonyms: 9G8, AAG3, SFRS7, Serine/arginine-rich splicing factor 7, Splicing factor 9G8,		
Splicing factor, arginine/serine-rich 7, SRSF7		
Background: Required for pre-mRNA splicing. Can also modulate alternative splicing in vitro.		
Represses the splicing of MAPT/Tau exon 10. May function as export adapter involved in		
mRNA nuclear export such as of histone H2A. Binds mRNA which is thought to be transferred		
to the NXF1-NXT1 heterodimer for export (TAP/NXF1 pathway), enhances NXF1-NXT1 RNA-		
binding activity. RNA-binding is semi-sequence specific.		
6432		
Q16629		
ELISA 1:500-1000		
IHC-P 1:200-400		
IHC-F 1:100-500		
IF(IHC-P) 1:50-200		
IF(IHC-F) 1:50-200		
IF(ICC) 1:50-200		
ICC 1:100-500		
For Research Use only		
Liquid		
1 μg/μL		
0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.		
ProClin		
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
handled by trained staff only.		
4 °C,-20 °C		
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
12 months		



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Formalin-fixed and paraffin embedded rat brain labeled with Anti-SFRS7/9G8 Polyclonal Antibody, Unconjugated at 1:200 followed by conjugation to the secondary antibody and DAB staining