antibodies -online.com





anti-TOR1B antibody (AA 281-336)



Go to Product page

()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL	
Target:	TOR1B	
Binding Specificity:	AA 281-336	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This TOR1B antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Torsin B
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Horse
Purification:	Purified by Protein A.

Target Details

|--|

Target Details

Abstract:	TOR1B Products	
Background:	Synonyms: DQ1, Tor1b, Torsin 1B, TOR1B, FKSG18, Torsin family 1 member B, TOR1B_HUMAN	
	Background: TorsinB is a 336 amino acid protein that localizes to the lumen of endoplasmic	
	reticulum and belongs to the Torsin family. Expressed in a variety of tissues with particularly	
	high levels in adult brain, TorsinB is thought to function as a molecular chaperone that assists	
	in the conformational folding of membrane and secreted proteins. Via its ability to regulate	
	protein folding, TorsinB plays a key role in postnatal developmental events and is essential for	
	proper neurological development. TorsinB contains two PKC (protein kinase C)	
	phosphorylations sites, one ATP binding domain and several conserved cysteines and shares	
	69 % amino acid identity with TorsinA, a related family member. Defects in the gene encoding	
	TorsinB are associated with torsion dystonia, an autosomal dominant neurological disorder	
	that is characterized by debilitating muscle contractions throughout the body.	
Application Details		
Application Notes:	WB 1:300-5000	
	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
	ICC 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	

Handling

Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months