# antibodies .- online.com







## anti-GLT8D2 antibody (AA 85-180)



$\sim$			
	N/6	1//r	$I \cap V$

Quantity:	100 μL
Target:	GLT8D2
Binding Specificity:	AA 85-180
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GLT8D2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human GLT8D2
Isotype:	lgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Horse,Rabbit
Purification:	Purified by Protein A.

#### Target Details

#### Target Details

Alternative Name:	Glt8d2 (GLT8D2 Products)	
Background:	Synonyms: GALA4A, GL8D2_HUMAN, GLT8D2, Glycosyltransferase 8 domain containing 2,	
	Glycosyltransferase 8 domain-containing protein 2, Gycosyltransferase.	
	Background: GLT8D2 (glycosyltransferase 8 domain-containing protein 2), also known as	
	GALA4A, is a 349 amino acid single-pass type II membrane protein. A member of the	
	glycosyltransferase 8 family, GLT8D2 is encoded by a gene that maps to human chromosome	
	12q23.3. Encoding over 1,100 genes within 132 million base pairs, chromosome 12 makes up	
	about 4.5 % of the human genome. A number of skeletal deformities are linked to chromosome	
	12, including hypochondrogenesis, achondrogenesis, Noonan syndrome, Kniest dysplasia and	
	trisomy 12p. Chromosome 12 is also home to a homeobox gene cluster, which encodes crucia	
	transcription factors for morphogenesis, as well as the natural killer complex gene cluster,	
	which encodes C-type lectin proteins that mediate the NK cell response to MHC I interaction.	
Gene ID:	83468	
Application Details		
Application Notes:	WB 1:300-5000	
	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	

### Handling

Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months