# antibodies -online.com





## anti-NHEDC2 antibody



Go to Product page

$\sim$				
	11/6	٦r١	/10	۱۸.

Quantity:	100 μL
Target:	NHEDC2
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NHEDC2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human NHEDC2
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

### **Target Details**

Target:	NHEDC2	
Alternative Name:	NHEDC2 (NHEDC2 Products)	
Background:	Synonyms: NHA2, Mitochondrial Na+/H+ exchanger NHA2, Mitochondrial sodium/hydroge exchanger NHA2, Na+/H+ exchanger like domain containing protein 2, Na+/H+ exchanger domain containing 2, NHE domain containing protein 2, NHE10, NHEDC 2, Sodium/hydrogen	

1	1:1	al a sala a toa	and the second		OLODO LILINANI
exchanger	IIKE	domain	containing	protein 2.	SL9B2_HUMAN.

Background: Na+/H+ exchangers (NHEs) catalyze the transport of Na+ in exchange for H+ across membranes in organisms and are required for numerous physiological processes.

NHEDC2 (Na+/H+ exchanger-like domain-containing protein 2), also known as NHA2, is a 537 amino acid mitochondrial protein. NHEDC2 is involved in organelle volume homeostasis by catalyzing the exchange of protons for Na+ and Li+ across the inner mitochondrial membrane. Found in red blood cells, NHEDC2 is required for bone resorption activity and osteoclast differentiation. As a multi-pass membrane protein, NHEDC2 is expressed as two isoforms produced by alternative splicing events.

Gene ID: 133308

Pathways: Proton Transport

#### **Application Details**

Application Notes: WB 1:300-5000

IHC-P 1:200-400

IF(IHC-P) 1:50-200

Restrictions: For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months