antibodies - online.com







anti-FAM98A antibody (AA 251-350)



| () | 1/0 | r\ /1 | 014 | |
|-----|-----|-------|-----|---|
| () | ve | I V I | -v | V |
| | | | | |

| Quantity: | 100 μL |
|----------------------|---|
| Target: | FAM98A |
| Binding Specificity: | AA 251-350 |
| Reactivity: | Human, Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This FAM98A antibody is un-conjugated |
| Application: | Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human FAM98A |
|-----------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse |
| Predicted Reactivity: | Rat,Cow,Sheep,Pig,Horse,Chicken |
| Purification: | Purified by Protein A. |

Target Details

| Target: | FAM98A | | |
|---------|--------|--|--|
|---------|--------|--|--|

Target Details

| Alternative Name: | FAM98A (FAM98A Products) | |
|---------------------|---|--|
| Background: | Synonyms: FAM 98A, Family with sequence similarity 98 member A, Hypothetical protein | |
| | LOC25940, LOC25940, Protein FAM98A, FA98A_HUMAN. | |
| | Background: Encoding more than 700 genes, chromosome 15 is made up of approximately 106 | |
| | million base pairs and is about 3 % of the human genome. Angelman and Prader-Willi | |
| | syndromes are associated with loss of function or deletion of genes in the 15q11-q13 region. Ir | |
| | the case of Angelman syndrome, this loss is due to inactivity of the maternal 15q11-q13 | |
| | encoded UBE3A gene in the brain by either chromosomal deletion or mutation. In cases of | |
| | Prader-Willi syndrome, there is a partial or complete deletion of this region from the paternal | |
| | copy of chromosome 15. Tay-Sachs disease is a lethal disorder associated with mutations of | |
| | the HEXA gene, which is encoded by chromosome 15. Marfan syndrome is associated with | |
| | chromosome 15 through the FBN1 gene. The FAM98 gene product has been provisionally | |
| | designated FAM98 pending further characterization. | |
| Gene ID: | 25940 | |
| Pathways: | SARS-CoV-2 Protein Interactome | |
| Application Details | | |
| Application Notes: | ELISA 1:500-1000 | |
| | FCM 1:20-100 | |
| | IHC-P 1:200-400 | |
| | IHC-F 1:100-500 | |
| | IF(IHC-P) 1:50-200 | |
| | IF(IHC-F) 1:50-200 | |
| | IF(ICC) 1:50-200 | |
| | ICC 1:100-500 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol. | |
| Preservative: | ProClin | |
| | | |

Handling

| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
|--------------------|--|
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. |
| Expiry Date: | 12 months |