antibodies -online.com





anti-MKRN2 antibody



Overview

Quantity:	100 μL
Target:	MKRN2
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MKRN2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human MKRN2/RNF62
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	MKRN2
Alternative Name:	RNF62 (MKRN2 Products)
Background:	Synonyms: HSPC070_ Makorin 2, Makorin ring finger protein 2, Makorin2, Makorin-2 MKRN 2, RING finger protein 62, RNF 62, RNF62, MKRN2_HUMAN, Probable E3 ubiquitin-protein ligase
	makorin-2, RING finger protein 62.

Background: The Makorins are a family of putative ribonucleoproteins containing two to four C3H zinc fingers that may confer RNA-binding. In addition, they contain a C3HC4 RING zinc finger that allows them to function as E3 ubiquitin ligases. Makorin-2, also known as RNF62, HSPC070 or MKRN2, is a widely expressed, evolutionarily conserved protein with four C3H-type zinc fingers (three at the N-terminus and one at the C-terminus), one RING-type zinc finger and a cysteine and histidine motif similar to that found in Makorin-1. In Xenopus, Makorin-2 functions, via PI 3-kinase/Akt signaling, as a negative regulator of neurogenesis. In humans, Makorin-2 is overexpressed in various cancer cell lines, suggesting a possible role of Makorin-2 in tumor progression. In addition, Makorin-2 is co-expressed with Raf-1 in the same tissues and cell lines.

Gene ID:

23609

Application Details

Application Notes: WB 1:300-5000

IHC-P 1:200-400

IF(IHC-P) 1:50-200

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months