antibodies -online.com





Datasheet for ABIN1385994

anti-APBB3 antibody (AA 401-486)



\sim		_		
(-, -)	\cap	\cup r \cap \cap	liiot	page
UU	w	IIOU	IUCL	vauc

()	ve	K\ /		A .
	\cup	1 V/	-	V۷

Quantity:	100 μL
Target:	APBB3
Binding Specificity:	AA 401-486
Reactivity:	Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This APBB3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), ELISA, Immunocytochemistry (ICC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human APBB3/FE65L2
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Cow,Sheep
Purification:	Purified by Protein A.

Target Details

Target:	APBB3		
---------	-------	--	--

Target Details

Alternative Name:	APBB3/FE65L2 (APBB3 Products)
Background:	Synonyms: Amyloid beta A4 precursor protein binding family B member 3, Amyloid beta A4
	precursor protein-binding family B member 3, amyloid beta precursor protein binding family B
	member 3, amyloid precursor interacting protein, Apbb3, APBB3_HUMAN, Fe65 like protein 2,
	FE65L2, Protein Fe65-like 2, SRA.
	Background: Fe65L2 is a 486 amino acid protein that contains one WW domain and two PID
	domains. Binding to the intracellular domain of the -Amyloid precursor protein, Fe65L2 is
	thought to modulate the internalization and, therefore, the accessibility and function of -
	Amyloid. Via its ability to control the intracellular accumulation of -Amyloid, Fe65L2 is though
	to play a role in the pathogenesis of Alzheimer's disease. Fe65L2 exists as four alternatively
	spliced isoforms designated isoform I, isoform II, isoform III and isoform IV. Fe65L2 interacts
	with Amyloid-like protein and is encoded by a gene located on human chromosome 5, which
	contains 181 million base pairs and comprises nearly 6 % of the human genome. Deletion of
	the p arm of chromosome 5 leads to Cri du chat syndrome, while deletion of the q arm or of
	chromosome 5 altogether is common in therapy-related acute myelogenous leukemias and
	myelodysplastic syndrome.
Gene ID:	10307
UniProt:	095704
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL

Handling

Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	
Expiry Date:	12 months	