antibodies -online.com





Datasheet for ABIN1386002 anti-Junctophilin 3 antibody (AA 181-280)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL
Target:	Junctophilin 3 (JPH3)
Binding Specificity:	AA 181-280
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Junctophilin 3 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Junctophilin 3
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken,Zebrafish
Purification:	Purified by Protein A.

Target Details

Target:	Junctophilin 3 (JPH3)
Alternative Name:	Junctophilin 3 (JPH3 Products)

Target Details

Background:

Synonyms: JP-3, JP3, JPH3, JPH3_HUMAN, Junctophilin3, Junctophilin type 3, Junctophilin-3, TNRC22, Trinucleotide repeat-containing gene 22 protein.

Background: Junctophilins are components of the junctional complexes between plasma membranes and endoplasmic or sarcoplasmic reticulums present in all excitable cells. Junctophilins contain a cytoplasmic domain which binds to the plasma membrane, as well as an ER/SR membrane spanning hydrophobic C-terminal segment. The three subtypes in this family are Junctophilin-1, -2 and -3. Junctophilin-1 is predominantly expressed in skeletal muscle, but is also expressed at low levels in heart. Junctophilin-2 is expressed in heart and skeletal muscle. Mutant mice lacking the Jph2 gene exhibit embryonic lethality and possess cardiac myocytes that express abnormal calcium transients. Junctophilin-3 is expressed in brain. The JPH3 alternatively spliced exon 2A has been suggested as a site for CTG repeat expansion leading to a Huntington disease-like autosomal dominant disorder.

Application Details

Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C

Handling

Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months