# antibodies -online.com





# anti-DCUN1D4 antibody (AA 161-270)



( )	11/0	K\ /	iew	1
	$\cup$	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	DCUN1D4
Binding Specificity:	AA 161-270
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DCUN1D4 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DCUN1D4
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken
Purification:	Purified by Protein A.

#### **Target Details**

|--|

# **Target Details**

- Target Details		
Alternative Name:	DCUN1D4 (DCUN1D4 Products)	
Background:	Synonyms: DCN1, defective in cullin neddylation 1, domain containing 4, DCN1 like protein 4,	
	DCUN1 domain containing protein 4, DCUN1D 4, KIAA0276, DCNL4_HUMAN.	
	Background: The DCN1-like protein family is comprised of Dcun1D1, Dcun1D2, Dcun1D3,	
	Dcun1D4 and Dcun1D5. The founding member, Dcun1D1, is involved in the malignant	
	transformation of squamous cell lineage.Dcun1D4, (defective in cullin neddylation protein 1-like	
	protein 4 or DCN1-like protein 4), also designated KIAA0276, exists as 2 isoforms as a result of	
	alternative splicing and contains one DCUN1 domain. The gene encoding Dcun1D4 maps to	
	chromosome 4, which houses nearly 6 % of the human genome and has the largest gene	
	deserts (regions of the genome with no protein encoding genes) of all of the human	
	chromosomes. Defects in some of the genes located on chromosome 4 are associated with	
	Huntington's disease, Ellis-van Creveld syndrome, methylmalonic acidemia and polycystic	
	kidney disease.	
Gene ID:	23142	
Application Details		
Application Notes:	WB 1:300-5000	
	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	

# Handling

Storage:	4 °C,-20 °C
Storage Comment: Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	
Expiry Date:	12 months