antibodies .- online.com







anti-SOX12 antibody (AA 51-150)



_			
\cup)ve	rvi	ew

Quantity:	100 μL
Target:	S0X12
Binding Specificity:	AA 51-150
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SOX12 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

- Toddet Details	
Immunogen:	KLH conjugated synthetic peptide derived from human SOX22
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Cow
Purification:	Purified by Protein A.
Target Details	

Target:	SOX12
Alternative Name:	SOX22/SOX12 (SOX12 Products)

Target Details

Background:

Synonyms: Homo sapiens SRY sex determining region Y box 12, Protein SOX-22, SOX 12, SOX 12 protein, SOX 22, SOX 22 protein, sox12, SOX12 protein, SOX12_HUMAN, SOX22 protein, SRY sex determining region Y box 12, SRY sex determining region Y box 22, SRY box 12, SRY box 22, SRY related HMG box gene 22, Transcription factor Sox-12.

Background: Sox genes comprise a family of genes that are related to the mammalian sex determining gene SRY. These genes similarly contain sequences that encode for the HMG-box domain, which is responsible for the sequence-specific DNA-binding activity. Sox genes encode putative transcriptional regulators implicated in the decision of cell fates during development and the control of diverse developmental processes. The highly complex group of Sox genes cluster at least 40 different loci that rapidly diverged in various animal lineages. At present, 30 Sox genes have been identified. Members of this family have been shown to be conserved during evolution and to play key roles during animal development. Some are involved in human diseases, including sex reversal.

Application Details

Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C

Handling

Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months