## .-online.com antibodies

## Datasheet for ABIN1386064 anti-RNF165 antibody



Overview

Quantity:	100 μL
Target:	RNF165
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RNF165 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human RNF165
Isotype:	lgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

## Target Details

Target:	RNF165
Alternative Name:	RNF165 (RNF165 Products)
Background:	Synonyms: ring finger protein 165, RN165_HUMAN.
	Background: The RING-type zinc finger motif is present in a number of viral and eukaryotic proteins and is made of a conserved cysteine-rich domain that is able to bind two zinc atoms.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1386064 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

	Proteins that contain this conserved domain are generally involved in protein-protein
	interactions and protein-DNA interactions. Specifically, they are thought to be involved in the
	ubiquitination pathway of protein degradation. RNF165 (RING finger protein 165) is a 346
	amino acid protein that contains one RING-type zinc finger. Via its RING-type zinc finger,
	RNF165 may play a role in transcriptional regulation and protein degradation events. The gene
	encoding RNF165 maps to human chromosome 18, which houses over 300 genes, including
	Trisomy 18, or Edwards syndrome, the second most common trisomy after Downs syndrome.
	Translocation between chromosome 18 and 14 is the most common translocation in cancers,
	and occurs in follicular lymphomas. Niemann-Pick disease, hereditary hemorrhagic
	telangiectasia and erythropoietic protoporphyria are also associated with chromosome 18.
Gene ID:	494470
Application Details	
Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN1386064 | 03/07/2024 | Copyright antibodies-online. All rights reserved.