antibodies -online.com





anti-FNTA antibody



Go to Product page

\sim					
()	VE	۲۱	/1	\triangle	Λ

Quantity:	100 μL	
Target:	FNTA	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This FNTA antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PGGT1A/FNTA	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified by Protein A.	

Target Details

Target:	FNTA
Alternative Name:	PGGT1A (FNTA Products)
Background:	Synonyms: CAAX farnesyltransferase alpha subunit, Farnesyl protein transferase alpha subunit, Farnesyltransferase CAAX box alpha, Farnesyltransferase, CAAX box, alpha, FPTA, FTase alpha,
	GGTase I alpha, PGGT1A, Protein farnesyltransferase/geranylgeranyltransferase type I alpha

subunit, Protein prenyltransferase alpha subunit repeat containing 2, PTAR2, Ras proteins prenyltransferase alpha, Ras proteins prenyltransferase subunit alpha, Type I protein geranyl geranyltransferase alpha subunit, FNTA_HUMAN.

Background: FNTA, also known as CAAX farnesyltransferase (FTase), attaches a farnesyl group from farnesyl pyrophosphate to cysteine residues at the fourth position from the C terminus of proteins that end in the so-called CAAX box, where C is cysteine, A is usually but not always an aliphatic amino acid, and X is typically methionine or serine. This type of posttranslational modification provides a mechanism for membrane localization of proteins that lack a transmembrane domain. This enzyme has the remarkable property of farnesylating peptides as short as four residues in length that conform to the CAAX consensus sequence. FNTA is also a specific cytoplasmic interactor of the transforming growth factor-beta and activin type I receptors. It is likely to be a key component of the signaling pathway which involves p21ras, an important substrate for farnesyltransferase.

Gene ID: 2339

Pathways: Response to Water Deprivation, Regulation of G-Protein Coupled Receptor Protein Signaling

Application Details

Application Notes: WB 1:300-5000

IHC-P 1:200-400

IF(IHC-P) 1:50-200

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

\vdash	land	lına
	iaria	шц

Expiry Date:

12 months