antibodies -online.com





anti-RNF146 antibody (AA 95-160)



Go to Product pag

\sim			
	N/P	r\/	i⊢₩

Quantity:	100 μL
Target:	RNF146
Binding Specificity:	AA 95-160
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RNF146 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human RNF146	
Isotype:	IgG	
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Pig, Horse, Chicken	
Purification:	Purified by Protein A.	

Target Details

Target:	RNF146
Alternative Name:	RNF146 (RNF146 Products)

Target Details

Background:

Synonyms: Dactylidin, dJ351K20.1, DKFZP434O1427, E3 ubiquitin-protein ligase rnf146, RING finger protein 146, RN146_HUMAN, RNF 146, Rnf146, RP3 351K20.1.

Background: The RING-type zinc finger motif is present in a number of viral and eukaryotic proteins and is made of a conserved cysteine-rich domain that is able to bind two zinc atoms. Proteins that contain this conserved domain are generally involved in the ubiquitination pathway of protein degradation. RNF146 (RING finger protein 146), also known as Dactylidin, is a 359 amino acid protein that contains one RING-type zinc finger and one WWE domain. Via its RING-type zinc finger, RNF146 may play a role in transcriptional regulation and protein degradation events. Defects in the gene encoding RNF146 are associated with Alzheimer's disease (AD) and may lead to a higher risk of breast cancer. Two isoforms of RNF146 exist due to alternative splicing events.

Application Details

Application Notes:	WB 1:300-5000

ELISA 1:500-1000

IHC-P 1:200-400

IHC-F 1:100-500

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

ICC 1:100-500

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

\vdash	land	lına
	iaria	шц

Expiry Date:

12 months