.-online.com antibodies

# Datasheet for ABIN1386149 anti-PCGF2 antibody (AA 170-210)

2 Images



#### Overview

Quantity:	100 µL
Target:	PCGF2
Binding Specificity:	AA 170-210
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PCGF2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

# Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human MEL18
Isotype:	lgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

### Target Details

Target:	PCGF2
Alternative Name:	MEL18 (PCGF2 Products)
Background:	Synonyms: MEL-18, RNF110, ZNF144, Polycomb group RING finger protein 2, DNA-binding

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN1386149 | 03/07/2024 | Copyright antibodies-online. All rights reserved. protein Mel-18, RING finger protein 110, Zinc finger protein 144, PCGF2, MEL18 Background: Transcriptional repressor. Binds specifically to the DNA sequence 5'-GACTNGACT-3'. Has tumor suppressor activity. May play a role in control of cell proliferation and/or neural cell development. Regulates proliferation of early T progenitor cells by maintaining expression of HES1. Also plays a role in antero-posterior specification of the axial skeleton and negative regulation of the self-renewal activity of hematopoietic stem cells. Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones, it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility. Is not functionally redundant with BMI1, unlike BMI1 does not stimulate the E3 ubiquitin-protein ligase activity in a reconstituted PRC1-like complex (By similarity).

Gene ID:	7703
UniProt:	P35227
Pathways:	Chromatin Binding

# Application Details

Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only


## Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN1386149 | 03/07/2024 | Copyright antibodies-online. All rights reserved. Expiry Date:

12 months

### Images



#### Immunohistochemistry

**Image 1.** Formalin-fixed and adjacent noncancerous of human colon tissue labeled with Anti-Mel18/ZNF144 Polyclonal Antibody, Unconjugated (ABIN1386149) at 1:200 followed by conjugation to the secondary antibody and DAB staining

#### Immunohistochemistry (Paraffin-embedded Sections)

**Image 2.** Formalin-fixed and adjacent noncancerous of human colon tissue labeled with Anti-Mel18/ZNF144 Polyclonal Antibody, Unconjugated at 1:200 followed by conjugation to the secondary antibody and DAB staining

