antibodies .- online.com







anti-MYLK antibody (AA 651-750)



Image



\sim						
	1//	Д	r۱	1	Θ 1	٨

Quantity:	100 μL	
Target:	MYLK	
Binding Specificity:	AA 651-750	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This MYLK antibody is un-conjugated	
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human MLCK
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human, Mouse, Dog, Cow, Sheep, Pig, Horse, Chicken, Guinea Pig
Purification:	Purified by Protein A.
T	
Target Details	

Target: MYLK


Target Details

Alternative Name:	MLCK (MYLK Products)
Background:	Synonyms: KRP, AAT7, MLCK, MLCK1, MYLK1, s mMLCK, MLCK108, MLCK210, MSTP083,
	Myosin light chain kinase, smooth muscle, Kinase-related protein, Telokin, MYLK
	Background: This gene, a muscle member of the immunoglobulin gene superfamily, encodes
	myosin light chain kinase which is a calcium/calmodulin dependent enzyme. This kinase
	phosphorylates myosin regulatory light chains to facilitate myosin interaction with actin
	filaments to produce contractile activity. This gene encodes both smooth muscle and
	nonmuscle isoforms. In addition, using a separate promoter in an intron in the 3' region, it
	encodes telokin, a small protein identical in sequence to the C-terminus of myosin light chain
	kinase, that is independently expressed in smooth muscle and functions to stabilize
	unphosphorylated myosin filaments. A pseudogene is located on the p arm of chromosome 3.
	Four transcript variants that produce four isoforms of the calcium/calmodulin dependent
	enzyme have been identified as well as two transcripts that produce two isoforms of telokin.
	Additional variants have been identified but lack full length transcripts. [provided by RefSeq, Ju
	2008]
Gene ID:	4638
UniProt:	Q15746
Application Details	
Application Notes:	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
	Liquid
Format:	
Format: Concentration:	1 μg/μL
	1 μg/μL 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	e Comment: Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	
Expiry Date:	12 months	

Images



Immunohistochemistry

Image 1. Formalin-fixed and paraffin embedded rat colon labeled with Anti-MLCK Polyclonal Antibody, Unconjugated (ABIN1386218) at 1:200, followed by conjugation to the secondary antibody and DAB staining