

Datasheet for ABIN1386324

anti-Transglutaminase 2 antibody (AA 351-450)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	Transglutaminase 2 (TGM2)
Binding Specificity:	AA 351-450
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Transglutaminase 2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Transglutaminase 2
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	Transglutaminase 2 (TGM2)
---------	---------------------------

Target Details

Alternative Name:	Transglutaminase 2 (TGM2 Products)
Background:	Synonyms: TG 2, TGC, TG2, TGase C, TGase H, TGase-2, TGase2, Tgasell, ALPHA SUBUNIT, C polypeptide, G alpha h, G protein alpha subunit Gh class, G[a]h, Gh CLASS G ALPHA h, GNAH, GNAH G PROTEIN, Guanine nucleotide binding protein H polypeptide, H POLYPEPTIDE, Protein glutamine gamma glutamyltransferase 2, Protein glutamine gamma glutamyltransferase, Protein-glutamine gamma-glutamyltransferase 2, TGC, TGC GUANINE NUCLEOTIDE BINDING PROTEIN, TGM 2, TGM2, TGM2_HUMAN, Tissue transglutaminase, Tissue type transglutaminase, Transglutaminase 2, Transglutaminase 2 C polypeptide, Transglutaminase C, Transglutaminase H, Transglutaminase-2, tTG, tTGas. Background: Terminally differentiating mammalian epidermal cells acquire an insoluble, 10 to 20 nm thick protein deposit on the intracellular surface of the plasma membrane known as the cross-linked cell envelope (CE). The CE is a component of the epidermis that is generated through formation of disulfide bonds and g-glutamyl-lysine isodipeptide bonds, which are formed by the action of transglutaminases (TGases). TGases are intercellularly localizing, Ca²⁺-dependent enzymes that catalyze the formation of isopeptide bonds by transferring an amine on to glutamyl residues, thereby cross-linking glutamine residues and lysine residues in substrate proteins. TGases influence numerous biological processes, including blood coagulation, epidermal differentiation, seminal fluid coagulation, fertilization, cell differentiation and apoptosis. Human keratinocyte transglutaminase (TGase1) is a membrane associated, 817 amino acid protein. Human tissue transglutaminase (TGase2) is an endothelial cell specific, 687 amino acid protein.
Pathways:	Tube Formation , Thromboxane A2 Receptor Signaling

Application Details

Application Notes:	WB 1:300-5000 ELISA 1:500-1000 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 ICC 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months