# antibodies .- online.com







## anti-FOLR2 antibody



( )	ve	K\ /		A .
	$\cup$	1 V/	Щ.	V۷

Quantity:	100 μL
Target:	FOLR2
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FOLR2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human Folate receptor
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

#### **Target Details**

Target:	FOLR2
Alternative Name:	FOLR2 (FOLR2 Products)
Background:	Synonyms: fetal/placental, Folbp 2, beta hFR, FBP, FBP/PL 1, FBP2, folate receptor 2 fetal,
	Folate receptor 2, Folate receptor, Folate receptor beta, Folate receptor, fetal/placental, Folbp2,
	Folr2, FOLR2_HUMAN, FR beta, FR P3, FR-beta, Placental folate binding protein, Placental

### **Target Details**

- arget betane	
	folate-binding protein.  Background: Folate is an essential vitamin that must be obtained from food intake through intestinal absorption in mammals. Folate and reduced folic acid derivatives bind to the folate receptor (FR) family, which mediates the endocytosis of 5-methyltetrahydofolate into the cell.
Pathways:	Dicarboxylic Acid Transport
Application Details	
Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months