# antibodies -online.com





## anti-LSAMP antibody (AA 51-150)



#### Overview

Quantity:	100 μL
Target:	LSAMP
Binding Specificity:	AA 51-150
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LSAMP antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC)

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human LSAMP
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Pig, Horse, Chicken, Rabbit
Purification:	Purified by Protein A.

### **Target Details**

Target:	LSAMP
Alternative Name:	LSAMP (LSAMP Products)

#### **Target Details**

#### Background:

Synonyms: IgLON family member 3, IGLON3, Lam, LAMP, Limbic system associated membrane protein, Limbic system associated membrane protein precursor, Limbic system-associated membrane protein, LSAMP, LSAMP\_HUMAN.

Background: LSAMP is a neuronal surface glycoprotein found in cortical and subcortical regions of the limbic system expressed on the surface of somata and proximal dendrites of neurons. It is a member of the immunoglobulin (Ig) superfamily and belongs to the IgLON subfamily of cell adhesion molecules. LSAMP contains three Ig domains and a glycosylphosphatidylinositol anchor. It is a highly conserved protein between rodents and humans (99 % sequence identity) and it is involved in the regulation of neurite outgrowth and mediation of proper circuit formation of limbic pathways. This suggests that LSAMP plays an important role in the development and function of the limbic system. In addition, LSAMP may function as a tumor suppressor in renal carcinomas.

#### **Application Details**

	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C

## Handling

Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months