antibodies -online.com







anti-NIPAL3 antibody (AA 1-100)



Go to Product page

()	11/	NI	/1/	≥W
	/ // 	۰۱ ا	// I E	- ¹ \/ \/

Quantity:	100 μL
Target:	NIPAL3
Binding Specificity:	AA 1-100
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NIPAL3 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human NIPAL3
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	NIPAL3	
rarget.	INII ALO	

Target Details

raiget Details	
Alternative Name:	NIPAL3 (NIPAL3 Products)
Background:	Synonyms: NPAL3, NPAL-3, NIPA like domain containing 3, NIPA like protein 3, RGD1563439,
	RP23-332E2.5, RP3-462023.3, NPAL3_HUMAN.
	Background: NIPAL3 is a 406 amino acid multi-pass membrane protein that belongs to the
	NIPA family and exists as three alternatively spliced isoforms. The gene that encodes NPAL3
	consists of approximately 57,229 bases and maps to human chromosome 1p36. Chromosome
	1 is the largest human chromosome spanning about 260 million base pairs and making up 8 $\%$
	of the human genome. There are about 3,000 genes on chromosome 1, and considering the
	great number of genes there are also a large number of diseases associated with chromosome
	1. Notably, the rare aging disease Hutchinson-Gilford progeria is associated with the LMNA
	gene which encodes lamin A. When defective, the LMNA gene product can build up in the
	nucleus and cause characteristic nuclear blebs. The MUTYH gene is located on chromosome 1
	and is partially responsible for familial adenomatous polyposis. Stickler syndrome, Parkinsons,
	Gaucher disease and Usher syndrome are also associated with chromosome 1.
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200

Handling	g

Restrictions:

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

IF(IHC-F) 1:50-200

For Research Use only

IF(ICC) 1:50-200

ICC 1:100-500

Handling

Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months