

Datasheet for ABIN1386444
anti-Trypsin Inhibitor antibody (AA 101-200)



[Go to Product page](#)

Overview

| | |
|----------------------|---|
| Quantity: | 100 µL |
| Target: | Trypsin Inhibitor |
| Binding Specificity: | AA 101-200 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This Trypsin Inhibitor antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC) |

Product Details

| | |
|-----------------------|---|
| Immunogen: | KLH conjugated synthetic peptide derived from human Trypsin Inhibitor |
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Predicted Reactivity: | Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken |
| Purification: | Purified by Protein A. |

Target Details

| | |
|---------|-------------------|
| Target: | Trypsin Inhibitor |
|---------|-------------------|

Target Details

Abstract: [Trypsin Inhibitor Products](#)

Background: Synonyms: Peptidase inhibitor 15, PI-15, 25 kDa trypsin inhibitor, p25TI, Cysteine-rich secretory protein 8, CRISP-8, SugarCrisp, PI15_HUMAN.

Background: Cysteine-rich secretory proteins (CRISPs) represent a family of evolutionarily conserved proteins which play a role in the innate immune system and are transcriptionally regulated by androgens in several tissues. CRISP-8 (Cysteine-rich secretory protein 8), also known as PI15 (Peptidase inhibitor 15), P25TI or SugarCrisp, is a 258 amino acid secreted protein that belongs to the CRISP family. Expressed at low levels in thyroid, prostate, salivary and mammary tissue, CRISP-8 functions as a serine protease inhibitor that exhibits weak inhibitory action against Trypsin, a serine protease found in the digestive system. In addition to its role as a protease inhibitor, CRISP-8 is secreted in neuroblastoma and glioblastoma cell lines, suggesting a role for CRISP-8 in tumor formation and metastasis within the central nervous system.

Application Details

Application Notes: WB 1:300-5000
ELISA 1:500-1000
IHC-P 1:200-400
IHC-F 1:100-500
IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200
ICC 1:100-500

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage: 4 °C,-20 °C

Storage Comment: Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Expiry Date: 12 months