# antibodies -online.com





# anti-EVC antibody (AA 251-350)



#### Overview

Quantity:	100 μL
Target:	EVC
Binding Specificity:	AA 251-350
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EVC antibody is un-conjugated
Application:	ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffinembedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC)

#### **Product Details**

Target:

Alternative Name:

**EVC** 

**EVC1 (EVC Products)** 

Immunogen:	KLH conjugated synthetic peptide derived from human EVC1	
Isotype:	IgG	
Predicted Reactivity:	Human	
Purification:	Purified by Protein A.	
Target Details		

#### Target Details

Dackground.	Bac	karo	und:
-------------	-----	------	------

Synonyms: Ellis van Creveld syndrome DWF 1, DWF1, Ellis van Creveld syndrome protein, Ellisvan Creveld syndrome, EVC, EVC1, EVC 1, EVC-1, EVCL, MGC105107, EVC\_HUMAN.

Background: EVC is an autosomal skeletal dysplasia caused by mutations in the EVC and EVC2 genes. Found in developing ribs, heart, kidney and lung, the EVC gene is responsible for normal development of the face, limbs, teeth and nails. The protein expressed by the EVC gene is an intracellular component of the hedgehog signal pathway that contains a leucine zipper and transmembrane domain. Defects in the EVC gene can lead to short-limb dwarfism, ectodermal dysplasia and cardiac anomalies such as irregular atrioventricular septum development.

Additionally, the EVC gene has been implicated in Weyers acrodental dysostosis, an autosomal dominant disease characterized by facial abnormalities and limb defects.

Pathways:

Hedgehog Signaling

### **Application Details**

Application Notes:

ELISA 1:500-1000

IHC-P 1:200-400

IHC-F 1:100-500

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

ICC 1:100-500

Restrictions:

For Research Use only

## Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months