antibodies - online.com





Datasheet for ABIN1386536

anti-SNAPIN antibody (AA 51-136)



Overview

Quantity:	100 μL
Target:	SNAPIN
Binding Specificity:	AA 51-136
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SNAPIN antibody is un-conjugated
Application:	ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffinembedded Sections) (IF (p)), Immunocytochemistry (ICC), Immunohistochemistry (Paraffinembedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SNAP25BP
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.
Target Details	

rarget Details

Target:	SNAPIN
Alternative Name:	SNAPIN/SNAP25BP (SNAPIN Products)

Target Details	
Background:	Synonyms: SNAP associated protein, SNAP-25-binding protein, SNAP-associated protein,
	SNAP25BP, SNAPAP, SNAPIN, SNAPN_HUMAN, SNARE associated protein snapin, SNARE-
	associated protein Snapin, Synaptosomal associated protein 25 binding protein, Synaptosomal-
	associated protein 25-binding protein, OTTHUMP00000035157.
	Background: The protein encoded by this gene is a coiled-coil-forming protein that associates
	with the SNARE (soluble N-ethylmaleimide-sensitive fusion protein attachment protein
	receptor) complex of proteins and the BLOC-1 (biogenesis of lysosome-related organelles)
	complex. Biochemical studies have identified additional binding partners. As part of the SNARE
	complex, it is required for vesicle docking and fusion and regulates neurotransmitter release.
	The BLOC-1 complex is required for the biogenesis of specialized organelles such as
	melanosomes and platelet dense granules. Mutations in gene products that form the BLOC-1
	complex have been identified in mouse strains that are models of Hermansky-Pudlak
	syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jun
	2012].
Gene ID:	23557
Pathways:	Synaptic Membrane, Synaptic Vesicle Exocytosis
Application Details	
Application Notes:	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months